arXiv:1211.6272v1 [gr-qc] 27 Nov 2012

Reconstruction of Scalar Potentials in Modified Gravity Models

A.Yu. Kamenshchik[1,2][*], **A. Tronconi**[1][†], **G. Venturi**[1][‡] and **S.Yu. Vernov**[3,4][§]

[1]Dipartimento di Fisica e Astronomia and INFN, Via Irnerio 46, 40126 Bologna, Italy,
[2]L.D. Landau Institute for Theoretical Physics of the Russian Academy of Sciences,
Kosygin str. 2, 119334 Moscow, Russia
[3]Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University,
Leninskie Gory 1, 119991, Moscow, Russia
[4]Instituto de Ciencias del Espacio (ICE/CSIC) and Institut d'Estudis Espacials de Catalunya,
Campus UAB, Fac. Ciències, Torre C5-Parell-2a pl., E-08193, Bellaterra (Barcelona), Spain

Abstract

We employ the superpotential technique for the reconstruction of cosmological models with a non-minimally coupled scalar field evolving on a spatially flat Friedmann–Robertson–Walker background. The key point in this method is that the Hubble parameter is considered as a function of the scalar field and this allows one to reconstruct the scalar field potential and determine the dynamics of the field itself, without a priori fixing the Hubble parameter as a function of time or of the scale factor. The scalar field potentials which lead to de Sitter or asymptotic de Sitter solutions, and those which reproduce the cosmological evolution given by Einstein-Hilbert action plus a barotropic perfect fluid, have been obtained.

1 Introduction

There are two possible ways of describing the accelerating expansion of the Universe [1]. One can assume that General Relativity is the correct theory of gravity and introduce a smoothly distributed, slowly varying cosmic fluid with negative pressure called dark energy [2]. Alternatively, one can modify the theory of gravity [3, 4, 5]. Some of these modified gravity models, for example, $F(R)$ gravity models, can then be mapped into general relativity with additional scalar fields by a suitable conformal transformation of the metric (see e.g. [5, 6]). It is well-known that scalar fields play an essential role in modern cosmology since they are possible candidates for the role of the inflaton field driving inflation in the early universe [7, 8] and of the dark energy substance [2]. The universe expansion history, unifying early-time inflation and late-time acceleration, can be realized in scalar-tensor gravity [9].

The modified gravity models, as well as cosmological models with scalar fields, often include some function which cannot be deduced from the fundamental theory. It is then a natural question to ask,

---

[*] Alexander.Kamenshchik@bo.infn.it
[†] Alessandro.Tronconi@bo.infn.it
[‡] Giovanni.Venturi@bo.infn.it
[§] svernov@theory.sinp.msu.ru, vernov@ieec.uab.es

why some specific form of such a function is chosen and what is the physical motivation for this choice. The technique of reconstruction has attracted the attention of researchers for a long time and has been developed for different cosmological models [10]–[35]. For example, the reconstruction of potentials for models with minimally coupled scalar fields has been studied in [10]–[20] and, in the two-field case, in [21, 22, 23], while a similar procedure for tachyon models was discussed in [24, 25, 26, 27]. The reconstruction of potentials for scalar fields non-minimally coupled to gravity was considered in [28, 29, 30]. We can also mention the reconstruction procedures in models with non-minimally coupled Yang–Mills fields [31], in $F(R)$ and Gauss–Bonnet gravity models [20, 32], in $F(T)$ models, with T being the torsion scalar [33], in the nonlocal gravity model [34] and its local formulation [35].

Gravity models with non-minimally coupled scalar fields, are actively studied in cosmology [36]–[43] (see also [5] and references therein). In particular the models with the Hilbert-Einstein term plus the term proportional to the Ricci scalar multiplied by the square of the scalar field were intensively studied in inflationary [44] and in quantum [45] cosmology.

Generally these models are described by the following action

$$S=\int d^4x\sqrt{-g}\left[U(\sigma)R-\frac{1}{2}g^{\mu\nu}\sigma_{,\mu}\sigma_{,\nu}+V(\sigma)\right], \tag{1}$$

where $U(\sigma)$ and $V(\sigma)$ are differentiable functions of the scalar field σ. We use the signature $(+,-,-,-)$ and g is the determinant of the metric tensor $g_{\mu\nu}$.

The reconstruction procedure for the induced gravity models ($U(\sigma)=\xi\sigma^2$ where ξ is a constant) has been proposed in [29]. In such a case it has been shown that one can linearize all the differential equations which should be solved in the reconstruction procedure to get the potential corresponding to a given cosmological evolution. This property allows one to obtain the explicit forms of potentials reproducing the dynamics of a flat Friedmann–Robertson–Walker (FRW) universe, driven by barotropic perfect fluids, by a Chaplygin gas and by a modified Chaplygin gas [29].

In this paper, we consider another reconstruction procedure for the models described by the above action (1). Such a method is similar to the Hamilton–Jacobi method (also known as the superpotential method) and is commonly applied to cosmological models with scalar fields [19, 21, 22, 23, 27] as well as in brane models [46]. The key point in this method is that the Hubble parameter is considered as a function of the scalar field σ.

The two methods described above supplement each other and together allow one to construct different cosmological models with some required properties. In particular, the reconstruction procedure illustrated in [29] is useful in order to obtain the potential and the explicit evolution of the scalar field when the Hubble parameter is given explicitly as a function of the cosmic time t or of the scale factor a. The superpotential method also allows one to do so (see Section 4), but for $U(\sigma)=\xi\sigma^2$ the procedure proposed in [29] is simpler. At the same time, however, the superpotential variant of the reconstruction procedure allows one to construct models and their exact solutions without any knowledge of the exact behavior of the Hubble parameter. For example, in Section 6 we describe the dynamics of a model with the Hubble parameter evolving to a nonzero constant value at late times. Such a method also allows one to calculate the function $U(\sigma)$, when the Hubble parameter is given as function of σ and the behavior of σ as a function of the cosmic time t (or the scale factor a) is known. Specifically, we assume that $\frac{d\sigma}{dt}=F(\sigma)$, where the function F is given. So, the superpotential procedure allows one to reconstruct not only the potential, but also either $U(\sigma)$, or the Hubble parameter, if the evolution of σ is given. Furthermore we can use this method to reconstruct scalar field potentials with some given property. For example, in Section 7 we find the conditions on the functions $U(\sigma)$ and $F(\sigma)$ for which the potential obtained is polynomial.

By using this method we find new exact solutions in the induced gravity case. We also applied the procedure to cosmological models with a non-minimally coupled scalar field described by (1) and

$U(\sigma)=\xi\sigma^2+J$, where J is a constant. In this physically important case, the method, proposed in [29], requires one to find solutions of second order nonlinear differential equations, whereas on using the superpotential method one only has to solve linear differential equations and first order autonomous nonlinear differential equations. Thus, the solutions can be always obtained at least in quadratures. One important case treated in detail is the reconstruction of de Sitter solutions. In such a case, the method proposed in [29] is still useful for arbitrary $U(\sigma)$ (see Section 3).

The paper is organized as follows. In Section 2, we review the basic equations for a gravity model with a non-minimally coupled scalar field in a FRW background and we describe the algorithm of the superpotential reconstruction procedure. In Section 3, we study the de Sitter solutions by using the methods proposed in [29] and in this paper and compare results. In Section 4, we find the non-minimally coupled models reproducing the dynamics of a flat FRW universe, driven by barotropic perfect fluids. In Section 5, we assume that the scalar field is proportional to the hyperbolic tangent (or cotangent) of the cosmic time and consider possible behaviors of the Hubble parameter and the corresponding potentials. In Section 6, we consider the induced gravity model and find solutions and potentials with a Hubble parameter which evolves towards a constant value at late times. In Section 7, we find conditions on the functions $U(\sigma)$ and $F(\sigma)$ sufficient to obtain a model with a polynomial potential. Finally Section 8 is devoted to the conclusions.

2 Cosmological models with non-minimally coupled scalar field

2.1 Homogeneous Equations

Consider the evolution of an homogeneous scalar field on a spatially flat FRW universe with the metric

$$ds^2=dt^2-a^2(t)\left(dx_1^2+dx_2^2+dx_3^2\right).$$

The Einstein equations derived from the variation of the action (1) have the following form[1] [29]:

$$6UH^2+6\dot{U}H=\frac{1}{2}\dot{\sigma}^2+V, \tag{2}$$

$$2U\left(2\dot{H}+3H^2\right)+4\dot{U}H+2\ddot{U}+\frac{1}{2}\dot{\sigma}^2-V=0, \tag{3}$$

where differentiation with respect to time t is denoted by a dot, and the Hubble parameter is the logarithmic derivative of the scale factor: $H=\dot{a}/a$. The variation of action (1) with respect to σ gives the Klein–Gordon equation:

$$\ddot{\sigma}+3H\dot{\sigma}+V_{,\sigma}=6\left(\dot{H}+2H^2\right)U_{,\sigma}\,, \tag{4}$$

where the subscript $_{,\sigma}$ indicates the derivative w.r.t. the scalar field σ. Combining Eqs. (2) and (3) we obtain:

$$4U\dot{H}-2\dot{U}H+2\ddot{U}+\dot{\sigma}^2=0. \tag{5}$$

This equation plays a key role in the reconstruction procedure.

---

[1] In [29], the authors have used a convenient choice of Newton's constant G, $8\pi G/3=1$, and dimensionless quantities. Alternatively, one can assume that G is a part of $U(\sigma)$ and do not fix the value of this constant.

2.2 The Hubble parameter as a function (superpotential) of the scalar field.

Let $H = Y(\sigma)$, then (5) takes the following form:

$$4UY_{,\sigma}\dot{\sigma} - 2YU_{,\sigma}\dot{\sigma} + 2U_{,\sigma\sigma}\dot{\sigma}^2 + 2U_{,\sigma}\ddot{\sigma} + \dot{\sigma}^2 = 0. \tag{6}$$

Let us introduce the function $F(\sigma)$ defined as

$$\dot{\sigma} = F(\sigma). \tag{7}$$

Substituting $\dot{\sigma}$ and $\ddot{\sigma} = F_{,\sigma}F$ into (6), we obtain:

$$4UY_{,\sigma} + 2(F_{,\sigma} - Y)U_{,\sigma} + (2U_{,\sigma\sigma} + 1)F = 0. \tag{8}$$

Equation (8) contains three functions. If two of them are given, then the third one can be found as the solution of a linear differential equation. Let us note that Eq. (8) is a first order differential equation both for Y and F.

If, for example, U and F are given, then Eq. (8) can be easily integrated to obtain $Y(\sigma)$:

$$Y(\sigma) = -\left(\int^{\sigma} d\tilde{\sigma}\frac{2F_{,\tilde{\sigma}}U_{,\tilde{\sigma}} + (2U_{,\tilde{\sigma}\tilde{\sigma}} + 1)F}{4U^{3/2}} + c_0\right)\sqrt{U} \tag{9}$$

where c_0 is an integration constant. In a similar fashion, for given $Y(\sigma)$ and $U(\sigma)$, we obtain $F(\sigma)$ by integrating (8):

$$F(\sigma) = \left(\int^{\sigma}\left[\frac{\frac{dU(\tilde{\sigma})}{d\tilde{\sigma}}Y(\tilde{\sigma}) - 2U(\tilde{\sigma})\frac{dY(\tilde{\sigma})}{d\tilde{\sigma}}}{\frac{dU(\tilde{\sigma})}{d\tilde{\sigma}}}e^{\Upsilon(\tilde{\sigma})}d\tilde{\sigma}\right] + \tilde{c}_0\right)e^{-\Upsilon(\sigma)}, \tag{10}$$

where

$$\Upsilon(\sigma) \equiv \frac{1}{2}\int^{\sigma}\frac{2\frac{d^2U}{d\tilde{\sigma}^2} + 1}{\frac{dU}{d\tilde{\sigma}}}d\tilde{\sigma}$$

and $\tilde{c}_0$ is an integration constant.

The potential $V(\sigma)$ can then be obtained from (2):

$$V(\sigma) = 6UY^2 + 6U_{,\sigma}FY - \frac{1}{2}F^2. \tag{11}$$

In order to find the time evolution $\sigma(t)$ we finally integrate Eq. (7), which can always be solved in quadratures.

The evolution of the Hubble parameter H and of the scalar field σ can be expressed either through the cosmic time t, or the scale factor a. In the latter case it is more convenient to adopt the dimensionless parameter $N \equiv \ln(a/a_0)$ as independent variable. In terms of N one has to re-express the time derivatives as

$$\frac{d}{dt} = H\frac{d}{dN} \quad \text{and} \quad \frac{d^2}{dt^2} = H^2\frac{d^2}{dN^2} + \frac{1}{2}\left(H^2\right)'\frac{d}{dN}, \tag{12}$$

where the prime denotes the derivative with respect to N. Equation (8) then takes the following form

$$2U(H^2)' - 2U'H^2 + 2U''H^2 + (H^2)'U' + (\sigma')^2H^2 = 0. \tag{13}$$

On defining the functions $W(\sigma)$ and $\tilde{F}(\sigma)$ as follows

$$W(\sigma) \equiv H^2 \tag{14}$$

and

$$\tilde{F}(\sigma) \equiv \sigma', \tag{15}$$

we find

$$(H^2)' = \frac{dH^2}{d\sigma}\frac{d\sigma}{dN} \equiv W_{,\sigma}\tilde{F}, \tag{16}$$

$$U' = U_{,\sigma}\tilde{F}, \qquad U'' = \left(U_{,\sigma\sigma}\tilde{F} + U_{,\sigma}\tilde{F}_{,\sigma}\right)\tilde{F} \tag{17}$$

and Eq. (13) can then be cast in the form:

$$2UW_{,\sigma} - 2U_{,\sigma}W + 2\left(U_{,\sigma\sigma}\tilde{F} + U_{,\sigma}\tilde{F}_{,\sigma}\right)W + W_{,\sigma}U_{,\sigma}\tilde{F} + \tilde{F}W = 0. \tag{18}$$

If, for example, $F(\sigma)$ and $U(\sigma)$ are given, the function $W(\sigma)$ can be determined by integrating:

$$\frac{W_{,\sigma}}{W} = \frac{\left[2U_{,\sigma} - 2\left(U_{,\sigma\sigma}\tilde{F} + U_{,\sigma}\tilde{F}_{,\sigma}\right) - \tilde{F}\right]}{\left(2U + U_{,\sigma}\tilde{F}\right)} \tag{19}$$

and finally the potential can be reconstructed by using

$$V = W\left[6U + 6U_{\sigma}\tilde{F} - \frac{1}{2}\tilde{F}^2\right]. \tag{20}$$

Equation (19) connects the functions W, U, and $\tilde{F}$ and is equivalent to Eq. (8). Indeed, on substituting $F = Y\tilde{F}$ into Eq. (8) we get

$$2\frac{Y_{,\sigma}}{Y} = \frac{\left[2U_{,\sigma} - 2\left(U_{,\sigma\sigma}\tilde{F} + U_{,\sigma}\tilde{F}_{,\sigma}\right) - \tilde{F}\right]}{\left(2U + U_{,\sigma}\tilde{F}\right)}. \tag{21}$$

The latter equation is equivalent to Eq. (19), because $W(\sigma) = Y^2(\sigma)$. We thus observe that the superpotential method allows one to unify the reconstruction procedures for the functions in (8) given as functions of t or of N. Indeed, one can start from Eq. (8) and find suitable functions Y, F and U satisfying it and, after that, it is trivial to get W and $\tilde{F}$.

2.3 Quadratic non-minimally coupled models

In this paper we consider

$$U(\sigma) = \xi\sigma^2 + C_1\sigma + J_0. \tag{22}$$

The case of induced gravity, for which the reconstruction procedure has been proposed in [29], corresponds to a particular choice of $U(\sigma)$ given by $C_1 = 0$ and $J_0 = 0$.

Note that the quadratic polynomial (22) can be rewritten as

$$U(\sigma) = \xi(\sigma - \sigma_0)^2 + J, \qquad \text{with} \quad \sigma_0 = -\frac{C_1}{2\xi}, \qquad J = J_0 - \frac{C_1^2}{4\xi}. \tag{23}$$

In this paper, we consider the case $\sigma_0 = 0$ (equivalent to $C_1 = 0$). Let us observe, however, that such a restriction does not lead to any loss of generality because any solution found with $\sigma_0 = 0$ can be extended to the $\sigma_0 \neq 0$ case simply through the change of variable $\sigma \to \sigma - \sigma_0$.

For $U(\sigma)$, given by (22) with $\sigma_0 = 0$, we obtain from (9) and (10):

$$Y(\sigma) = -\left[\int^{\sigma} \frac{4\xi\tilde{\sigma}F_{,\tilde{\sigma}} + (4\xi+1)F}{4(\xi\tilde{\sigma}^2 + J)^{3/2}}\, d\tilde{\sigma} + c_0\right]\sqrt{\xi\sigma^2 + J}\,, \tag{24}$$

$$F(\sigma) = \left[\int^{\sigma} \left[\tilde{\sigma}Y - \left(\tilde{\sigma}^2 + \frac{J}{\xi}\right)Y_{,\tilde{\sigma}}\right]\tilde{\sigma}^{1/(4\xi)}\, d\tilde{\sigma} + \tilde{c}_0\right]\sigma^{-1-1/(4\xi)}\,. \tag{25}$$

Let us note that the behavior of the Hubble parameter can be very different in a model with zero and nonzero J. To illustrate this point let us consider the example with an arbitrary linear function $F(\sigma)$:

$$F = f_1\sigma + f_0, \tag{26}$$

where f_i are constants. From Eq. (24), we get:

$$Y_J(\sigma) = \frac{f_1}{4\xi} + 2f_1 - \frac{(4\xi+1)f_0}{4J}\sigma + c_0\sqrt{\xi\sigma^2 + J}\,, \quad J \neq 0, \tag{27}$$

$$Y_0(\sigma) = \frac{f_1}{4\xi} + 2f_1 + \frac{(4\xi+1)f_0}{8\xi\sigma} + c_0\sigma\,, \quad J = 0. \tag{28}$$

Therefore, if $f_0 \neq 0$ and $\xi \neq -1/4$, then $Y_0(\sigma)$ is not a limit of $Y_J(\sigma)$ at $J \to 0$.

3 Models with de Sitter solutions

In this section we shall consider the general form of the potential $V(\sigma)$, which leads to the existence of the de Sitter solution

$$H = H_0 = const \tag{29}$$

for an arbitrary non-minimal coupling $U(\sigma)$. In this case, it is convenient to apply the technique developed in [29].

First of all let us notice, that if the scalar field σ does not depend on time t (or, in other words, on the scale factor a) then all the considerations presented in the end of the second section of paper [29] are still valid and we obtain immediately the de Sitter solution for the model with the potential (see Eq. (31) of the above cited paper)

$$V(\sigma) = V_0U^2(\sigma), \tag{30}$$

where V_0 is an arbitrary positive constant. The constant value of the Hubble parameter is then (see Eq. (2))

$$H_0 = \pm\sqrt{\frac{V}{6U}} = \pm\sqrt{\frac{V_0U}{6}}. \tag{31}$$

For the case of a time-dependent scalar field σ, we use Eq. (19) from paper [29] and we rewrite it in terms of $N \equiv \ln a/a_0$

$$\sigma'' - \sigma' + \sigma'\frac{H'}{H} + (\sigma')^2\left(\frac{1 + 2\frac{d^2U}{d\sigma^2}}{2\frac{dU}{d\sigma}}\right) + \frac{2UH'}{\frac{dU}{d\sigma}H} = 0. \tag{32}$$

For the de Sitter case, this equation reduces to

$$\sigma'' - \sigma' = -\left(\frac{1+2\frac{d^2U}{d\sigma^2}}{2\frac{dU}{d\sigma}}\right)(\sigma')^2. \tag{33}$$

Using $\frac{d}{dN} = \sigma'\frac{d}{d\sigma}$, after some algebra, one can integrate both sides of Eq. (33) and obtain:

$$\ln\sigma' - N = -\int^{\sigma} d\tilde{\sigma}\left(\frac{1+2\frac{d^2U}{d\tilde{\sigma}^2}}{2\frac{dU}{d\tilde{\sigma}}}\right) + \tilde{B}_0 = -\int^{\sigma}\frac{d\tilde{\sigma}}{2\frac{dU}{d\tilde{\sigma}}} - \ln\left(\frac{dU}{d\sigma}\right) + \tilde{B}_0. \tag{34}$$

where $\tilde{B}_0$ is an integration constant.

We now exponentiate and integrate the above equation to find $\exp(N)$ as a function of the scalar field σ:

$$\exp(N) = \frac{1}{B_0}\int^{\sigma} d\sigma_*\frac{dU}{d\sigma_*}\exp\left(\frac{1}{2}\int^{\sigma_*} d\tilde{\sigma}\frac{1}{\frac{dU}{d\tilde{\sigma}}}\right) + B_1, \tag{35}$$

where $B_0 = \exp(\tilde{B}_0)$ and B_1 is an arbitrary constant.

Consequently we obtain σ' as a function of σ:

$$\sigma' = \frac{\exp\left(-\frac{1}{2}\int^{\sigma} d\tilde{\sigma}\frac{1}{\frac{dU}{d\tilde{\sigma}}}\right)}{\frac{dU}{d\sigma}}\left[B_1B_0 + \int^{\sigma} d\sigma_*\frac{dU}{d\sigma_*}\exp\left(\frac{1}{2}\int^{\sigma_*} d\tilde{\sigma}\frac{1}{\frac{dU}{d\tilde{\sigma}}}\right)\right] \tag{36}$$

and the corresponding potential can be found by substituting (36) and (29) into Eq. (20).

For the particular case:

$$U(\sigma) = \xi\sigma^2 + J, \tag{37}$$

we obtain

$$\sigma' = \frac{4\xi}{1+8\xi}\sigma + \frac{B_1B_0}{2\xi}\sigma^{-\frac{1+4\xi}{4\xi}}. \tag{38}$$

Substituting (38) into (2) we get

$$V = H_0^2\left[6J + \frac{2(3+32\xi)(1+12\xi)\xi}{(8\xi+1)^2}\sigma^2 + \frac{4B_1B_0(1+12\xi)}{1+8\xi}\sigma^{-\frac{1}{4\xi}} - \frac{B_1^2B_0^2}{8\xi^2}\sigma^{-\frac{4\xi+1}{2\xi}}\right]. \tag{39}$$

Let us apply the superpotential method to the same model with the de Sitter evolution. Using (10), we obtain

$$F(\sigma) = \left[\int^{\sigma} H_0 e^{\Upsilon} d\tilde{\sigma} + \tilde{c}_0\right] e^{-\Upsilon} \tag{40}$$

and, in particular, for U, given by (37),

$$F(\sigma) = \frac{4\xi H_0}{8\xi+1}\sigma + \tilde{c}_0\sigma^{-1-1/(4\xi)}. \tag{41}$$

Therefore,

$$\sigma' = \tilde{F}(\sigma) = \frac{F(\sigma)}{Y(\sigma)} = \frac{4\xi}{8\xi+1}\sigma + \frac{\tilde{c}_0}{H_0}\sigma^{-1-1/(4\xi)}. \tag{42}$$

This formula coincides with (38), if we choose $\tilde{c}_0 = H_0 B_1 B_0/(2\xi)$, so we conclude that the potential can be reconstructed with both the procedures.

Solving Eq. (7), we obtain

$$\sigma(t) = \left[\sigma_0 e^{H_0 t} + \frac{\tilde{c}_0(8\xi+1)}{H_0\xi}\right]^{4\xi/(8\xi+1)}, \tag{43}$$

where σ_0 is an arbitrary constant.

For the case $\tilde{c}_0 = 0$, $F(\sigma)$ is a linear function. The potential V is the following quadratic polynomial:

$$V = 2H_0^2\left[3J + \frac{(3+32\xi)(1+12\xi)\xi}{(8\xi+1)^2}\sigma^2\right]. \tag{44}$$

For $\xi = -1/12$ and $\xi = -3/32$ the potential V is a constant.

The case $\xi = -1/8$ should be considered separately. In this case, using Eq. (10), we get:

$$F(\sigma) = \sigma H_0 \ln\left(\frac{\sigma}{\sigma_0}\right), \tag{45}$$

where σ_0 is an integration constant, and the corresponding potential has the following form:

$$V = \frac{H_0^2}{4}\left[24J - \frac{\sigma^2}{2}\left(\ln\left(\frac{\sigma}{\sigma_0}\right)^2 + 3 + \sqrt{3}\right)\left(\ln\left(\frac{\sigma}{\sigma_0}\right)^2 + 3 - \sqrt{3}\right)\right].$$

The scalar field evolution is given by

$$\sigma(t) = \sigma_0 \exp\left[e^{H_0(t-t_0)}\right]. \tag{46}$$

It is easy to see that in the case of induced gravity ($J = 0$) the expressions (39) and (44) can be derived from the expression (51) in the paper [29] provided we identify $c_1 = -\frac{B_1 B_0(1+8\xi)}{8\xi^2}$ and $\gamma = 2\xi$. Also the particular cases $\gamma = -1/6$ (conformal coupling) and $\gamma = -1/4$ corresponding to $\xi = -1/12$ and $\xi = -1/8$ respectively were considered in [29].

4 Power-law Solutions

In general relativity with a barotropic perfect fluid, the Hubble parameter evolves as

$$H^2(a) = H_0^2\left(\frac{a}{a_0}\right)^{-3(w+1)} = H_0^2 e^{nN}, \tag{47}$$

where the constant w is the equation-of-state parameter of the fluid and $n \equiv -3(w+1)$. Let us note that the Hubble parameter as a function of time is $H(t) = -\frac{2}{nt} = \frac{2}{3(w+1)t}$. Power-law solutions for induced gravity models ($J = 0$) and the corresponding potentials have been studied in [29]. Let us note that, in the case with $J \neq 0$, the method employed in [29] leads to a differential equation which cannot be solved analytically. For such a case it is convenient to adopt the superpotential technique for reconstruction.

Let us start from Eq. (19) and observe that the power-law dependence of H^2 on the scale factor gives $n = (H^2)'/H^2$, or equivalently

$$n = \frac{W_{,\sigma}\tilde{F}}{W}. \tag{48}$$

Then Eq. (19) becomes

$$\frac{n}{\tilde{F}} = \frac{\left[2U_{,\sigma} - 2\left(U_{,\sigma\sigma}\tilde{F} + U_{,\sigma}\tilde{F}_{,\sigma}\right) - \tilde{F}\right]}{\left(2U + U_{,\sigma}\tilde{F}\right)}\,. \tag{49}$$

After substituting (37), Eq. (49) can be recast in the following form:

$$4\xi\sigma\tilde{F}_{,\sigma}\tilde{F} + 2\xi\sigma(n-2)\tilde{F} + (4\xi+1)\tilde{F}^2 + 2nJ + 2n\xi\sigma^2 = 0. \tag{50}$$

In particular, for $n = 2$, Eq. (50) is a linear differential equation for $\tilde{F}^2$ which can be solved exactly leading to the following solution:

$$\tilde{F}(\sigma) = \pm\sqrt{-4\frac{(4\xi+1)\xi\sigma^2 + (8\xi+1)J}{(4\xi+1)(8\xi+1)} + c_1\sigma^{-(4\xi+1)/(2\xi)}} \tag{51}$$

For this solution from Eq. (48), we get

$$W(\sigma) = W_0 e^{\int^{\sigma}\frac{n}{\tilde{F}(\tilde{\sigma})}d\tilde{\sigma}}. \tag{52}$$

For the values of ξ leading to a singular expression (51), Eq. (50) has the following general solution:

- for $\xi = -1/4$

$$\tilde{F}(\sigma) = \pm\sqrt{8J\ln(\frac{\sigma}{\sigma_0}) - \sigma^2}, \tag{53}$$

- for $\xi = -1/8$

$$\tilde{F}(\sigma) = \pm\sqrt{-2\sigma^2\ln(\frac{\sigma}{\sigma_0}) - 8J} \tag{54}$$

where σ_0 is the integration constant.

It is impossible to solve Eq. (50) by the separation of σ and $\tilde{F}$ for the case of an arbitrary n. Hence we look for particular solutions of Eq. (50) by making a suitable ansatz for $\tilde{F}$ as a function of σ and then algebraically fixing the free coefficients of the model. We choose the ansatz

$$\tilde{F}(\sigma) = A_0 + A_1\sigma\,, \tag{55}$$

by which Eq. (50) reduces to a quadratic polynomial of σ, which can be solved by requiring that the coefficients of the diverse powers of σ are zero. Let us note that the above ansatz (55) when $A_1 \neq 0$ corresponds to the following evolution for the scalar field in terms of N

$$\sigma(N) = B\exp\left(A_1 N\right) - \frac{A_0}{A_1} \tag{56}$$

where B is an integration constant. It is then straightforward to obtain the evolution $\sigma(t)$ by integrating:

$$dN = \frac{dN}{dt}dt = Hdt = -\frac{2}{n\,t}dt \Rightarrow N = \ln\left(\frac{t}{t_0}\right)^{-2/n}. \tag{57}$$

and substituting into Eq. (56). Equation (50) gives three constraints on the 5 parameters of the model: A_0, A_1, n, ξ and J. If we assume that the function $U(\sigma)$ is given by (37), or in other words that the parameters ξ and J are fixed, we then find the following expressions for the remaining three parameters:

$$\begin{aligned} A_0^2 &= \frac{2\left[-14\xi - 44\xi^2 - 1 \pm (6\xi+1)\sqrt{52\xi^2+16\xi+1}\right] J}{(4\xi+1)^2\xi}, \\ A_1 &= \frac{-36\xi^2 - 12\xi - 1 \pm (6\xi+1)\sqrt{52\xi^2+16\xi+1}}{(6\xi+1)(4\xi+1)}, \\ n &= \frac{14\xi + 44\xi^2 + 1 \mp (6\xi+1)\sqrt{52\xi^2+16\xi+1}}{(4\xi+1)\xi}. \end{aligned} \tag{58}$$

These solutions exist only if $\xi \neq -1/4$ and $\xi \neq -1/6$. The parameters n and A_1 are real if and only if $\xi < (-4-\sqrt{3})/26$ or $\xi > (-4+\sqrt{3})/26$. Let us note that $A_0 = 0$ if and only if $J = 0$.

On the other hand, if we assume, that n is given and seek both the functions F and U then Eq. (50) leads to two nontrivial solutions with A_0 unconstrained and:

$$J_{1,2} = A_0^2 \frac{-4n - \alpha \pm (n-2)\sqrt{\alpha}}{4n\,(\alpha - 2n)} \tag{59}$$

$$\xi_{1,2} = \frac{8n - \alpha \mp (n-2)\sqrt{\alpha}}{8\,(\alpha - 2n)} \tag{60}$$

$$A_{1;1,2} = \frac{2 - n \mp \sqrt{\alpha}}{4} \tag{61}$$

with $\alpha = 4 - 20n + n^2$ and where A_0 and $-A_0$ correspond to the same values for the other parameters. These solutions describe a Universe evolving according to (47) due to the presence of a non-minimally coupled scalar field with the coupling to the Ricci scalar given by (60). Note that α is non negative and expressions (59)–(61) are real provided $n \geq 10 + 4\sqrt{6} \simeq 19.8$ or $n \leq 10 - 4\sqrt{6} \simeq 0.2$. Negative values of n correspond to a decreasing H in an expanding universe and describe relevant cosmological evolutions. The limit $n \to 0$ gives the de Sitter solution and the solutions with small, positive n still describe interesting cosmological evolutions corresponding to a super accelerated era (the fact that the Universe is currently undergoing a phase of super accelerated expansion is not excluded by supernovae observations [48]).

Equation (50) also has 4 more solutions corresponding either to the induced gravity or the general relativity framework. We find induced gravity solutions (with $J = 0$) having $A_0 = 0$, ξ unconstrained and

$$A_{1;3,4} = \frac{-(n-2)\xi \pm \sqrt{\xi\,(\alpha\,\xi - 2n)}}{1 + 8\xi} \tag{62}$$

or $\xi_5 = -1/8$ and $A_{1;5} = n/\,(2-n)$. In these cases the scalar field evolution is given by $\sigma(N) = B\exp A_1 N$. The general relativity solution is finally for $A_1 = 0$, $\xi = 0$, $J = -A_0^2/(2n)$ and a scalar field evolution given by $\sigma = A_0 N + B$ where B is an arbitrary constant. The constant A_0 is physically constrained by the value of the Planck mass in the Einstein–Hilbert action.

Eq. (48) can be solved for W with $\tilde{F}$ given by the ansatz (55):

$$W = W_0 \exp\left[\int^{\sigma} \frac{n}{\tilde{F}} d\tilde{\sigma}\right] = W_0 \exp\left[\int^{\sigma} \frac{n}{A_0 + A_1\tilde{\sigma}} d\tilde{\sigma}\right] = W_0\,(A_0 + A_1\sigma)^{\frac{n}{A_1}}\,. \tag{63}$$

where W_0 is an integration constant.

On using Eq. (20), we obtain the corresponding potential

$$V = W_0 \left(A_0 + A_1\sigma\right)^{\frac{n}{A_1}} \left[6\left(J + \xi\sigma^2\right) + 12\xi\sigma\left(A_0 + A_1\sigma\right) - \frac{\left(A_0 + A_1\sigma\right)^2}{2}\right] \tag{64}$$

with J, ξ and A_1 given by (59)–(61) respectively. In the induced gravity case (solutions 3 and 4) the potential takes the following form:

$$V_{3,4} = \tilde{W}_0 \sigma^{\frac{n}{A_{1;3,4}}+2} = \tilde{W}_0 \sigma^{\frac{(6-n)\xi \pm \sqrt{\xi(\alpha\xi-2n)}}{2\xi}} \tag{65}$$

where we absorbed all the parameters and the integration constant W_0 into $\tilde{W}_0$. Correspondingly the evolution of the scalar field is given by

$$\sigma_{3,4}(N) = B \exp\left[\frac{-(n-2)\xi \pm \sqrt{\xi\left(\alpha\,\xi - 2n\right)}}{1+8\xi} N\right] . \tag{66}$$

These two solution where already found in [29]. In the $n \to 0$ (and $\alpha \to 4$) limit one then obtains the de Sitter solutions (see [29]) with

$$V_3^{(dS)} = \tilde{W}_0 \sigma^2 \,, \quad V_4^{(dS)} = \tilde{W}_0 \sigma^4 \tag{67}$$

and

$$\sigma_3^{(dS)}(N) = B \exp\left[\frac{4\xi}{1+8\xi} N\right] \,, \quad \sigma_4^{(dS)}(N) = B. \tag{68}$$

If we consider the induced gravity solution left, with $\xi_5 = -1/8$, we are led to the potential

$$V_5 = \tilde{W}_0 \sigma^{4-n} \tag{69}$$

and the evolution of the scalar field is given by

$$\sigma_5(N) = B \exp\left[\frac{n}{2-n} N\right] . \tag{70}$$

Let us note that such a solution cannot be obtained from the solutions 1-4 with an appropriate limit procedure.

The general relativity solution has

$$W = W_0 \exp\left(\frac{n}{A_0}\sigma\right) \tag{71}$$

and a corresponding potential of the form

$$V = \tilde{W}_0 \exp\left(\frac{n}{A_0}\sigma\right) \tag{72}$$

which is the well-know exponential potential associated with power-law expansion.

The solutions 1 and 2 deserve some more discussion. In the limit $J_{1,2} \propto A_0 \to 0$ one obtains induced gravity with the non-minimal coupling fixed at $\xi = \xi_{1,2}$. In such a case the potential takes the form

$$V_{1,2} = \tilde{W}_0 \sigma^{\frac{10-n\pm\sqrt{\alpha}}{4}} \tag{73}$$

and the scalar fields evolves as

$$\sigma_{1,2}(N) = B \exp\left[\frac{2-n\mp\sqrt{\alpha}}{4}\right] . \tag{74}$$

Note that the solution 1, in the induced gravity limit, is the solution 4 with $\xi = \xi_1$ and the solution 2, in the same limit, is the solution 3 with $\xi = \xi_2$. Thus, when $J = 0$, the solutions 1 and 2 are particular cases of the solutions 3 and 4.

5 Solutions with the hyperbolic tangent

Let us construct cosmological models, when the Hubble parameter is a function of the hyperbolic tangent. Such solutions are popular in cosmology, because they can describe a bounce [47] and late-time acceleration [17, 21] having de Sitter/anti-de Sitter attractors both in the past and in the future. Such models can be reconstructed, for example, starting with the following ansatz for the scalar field evolution

$$\sigma(t) = A\tanh\left[\omega(t-t_0)\right], \tag{75}$$

where A, ω and t_0 are constants. Note that t_0 can be complex, so the parametrization (75) includes the functions $\sigma(t) = A\coth\left[\omega(t-t_0)\right]$ as well. For such functions Eq. (7) takes the form

$$F(\sigma) = \omega\left(A - \frac{1}{A}\sigma^2\right). \tag{76}$$

If $U(\sigma) = \xi\sigma^2$, then after substituting $F(\sigma)$, given by (76), into formula (24) we can integrate the resulting equation to obtain:

$$Y(\sigma) = \frac{(4\xi+1)\omega A}{8\xi\sigma} + \frac{(12\xi+1)\omega}{4\xi A}\sigma\ln\left(\frac{\sigma}{\sigma_0}\right), \tag{77}$$

where σ_0 is an arbitrary constant. The function $Y(\sigma)$ has the simplest form for $\xi = -1/12$:

$$Y(\sigma) = c_0\sigma - \frac{\omega A}{\sigma}. \tag{78}$$

The corresponding potential is the fourth degree monomial:

$$V = -\frac{(\omega - Ac_0)^2}{2A^2}\sigma^4. \tag{79}$$

For $c_0 = 0$, we obtain the following evolution for the Hubble parameter:

$$H = -\,\omega\coth[\omega(t-t_0)] \tag{80}$$

and conversely we get $H = -\omega\tanh[\omega(t-t_0)]$, when $\sigma(t) = A\coth[\omega(t-t_0)]$.

In the more general case with $U(\sigma) = -\sigma^2/12 + J$ we get:

$$Y = c_0\sqrt{12J - \sigma^2} - \frac{\omega A}{6J}\sigma. \tag{81}$$

For $c_0 = 0$, the Hubble parameter is proportional to σ and the potential has following polynomial structure:

$$V = -\,\frac{(6J+A^2)^2\omega^2}{72J^2A^2}\sigma^4 + \left(\frac{A^2}{3J}+1\right)\omega^2\sigma^2 - \frac{1}{2}\omega^2A^2. \tag{82}$$

Another way to get the desired Hubble parameter evolution (with constant H attractors in the past and in the future), is to express H as a function of σ evolving as (75) and then solve the Eq. (8) as a differential equation for $U(\sigma)$. On assuming

$$H = Y(\sigma) = B - C\sigma, \tag{83}$$

where B and C are constants, and then substituting (76) and (83) into Eq. (8), we get the following second order linear differential equation for $U(\sigma)$:

$$2\Omega(A^2-\sigma^2)U_{,\sigma\sigma} + 2\left[(C-2\Omega)\sigma - B\right]U_{,\sigma} - 4CU + (A^2-\sigma^2)\Omega = 0, \tag{84}$$

where we set $\omega = \Omega A$.

A particular solution of this equation is the second degree polynomial

$$U(\sigma) = -\frac{1}{12}\sigma^2 + \frac{B}{6(2\Omega + C)}\sigma + \frac{2A^2C\Omega + 4A^2\Omega^2 - B^2}{12(2\Omega + C)C}, \tag{85}$$

having the form (22). For any values of A, B, C and Ω, except $\Omega = -C$, the potential $V(\sigma)$ is a fourth degree polynomial with a coefficient of the high-order degree monomial equal to $-(\Omega + C)^2/2$.

Let us consider, for example, the particular set of parameters $B = A$ and $C = 1$. We get

$$U(\sigma) = -\frac{1}{12}\sigma^2 + \frac{A}{6(2\Omega + 1)}\sigma + \frac{(2\Omega + 4\Omega^2 - 1)A^2}{12(2\Omega + 1)}, \tag{86}$$

hence, $U(\sigma) = 0$ at

$$\sigma_{1,2} = A\frac{1 \pm 2\sqrt{2\Omega^2 + 2\Omega^3}}{2\Omega + 1}.$$

It is easy to see that $U(A) > 0$ for all $\Omega > 0$, so, if we choose as a solution $\sigma(t) = A\tanh(\omega t)$, then $U(t)$ is positive at late times if $A > 0$. We further observe that (see Fig. 1), when $\Omega = 1$, $U(\sigma(t)) \geq 0$ at any time because $-A \leqslant \sigma(t) \leqslant A$. For $\Omega = 1$, we get



Figure 1: The function $U(\sigma)$, given by (86), (left) and the potential $V(\sigma)$, given by (87), (right) at $A = 1$, $B = 1$, $C = 1$ and $\Omega = 1$.

$$V(\sigma) = -2\sigma^4 + \frac{8}{3}A\sigma^3 + \frac{4}{3}A^2\sigma^2 - \frac{8}{3}A^3\sigma + \frac{2}{3}A^4 = -\frac{2}{3}(\sigma + A)(3\sigma - A)(\sigma - A)^2 \tag{87}$$

and, by the change of variable $\tilde{\sigma} = \sigma - A/3$, the expression (86) becomes

$$U(\tilde{\sigma}) = -\frac{1}{12}\tilde{\sigma}^2 + \frac{4}{27}A^2, \tag{88}$$

and takes the form of (37) with $\xi = -1/12$. In terms of $\tilde{\sigma}$, we finally get

$$Y(\tilde{\sigma}) = \frac{2}{3}A - \tilde{\sigma}, \qquad F(\tilde{\sigma}) = \frac{(2A - 3\tilde{\sigma})(4A + 3\tilde{\sigma})}{9}. \tag{89}$$

So, we found a model with exact solutions and $U(\sigma)$ in the form $U(\sigma)=\xi\sigma^2+J$. We note that if we consider general values of A, C, and ω still the particular solution (85) is for $\xi=-1/12$. This value of ξ corresponds to the case of the conformal coupling [29].

6 Non-monotonic behavior of the Hubble Parameter in the Case of Induced Gravity

In the previous sections we considered the case of $U(\sigma)$, specified by (37). The case of induced gravity ($J=0$) has been considered in detail in [29], where another method of reconstruction has been used. In this section, we find new induced gravity cosmological models with exact solutions on using the superpotential method. We demonstrate that the same evolution $\sigma(t)$ leads to exactly solvable models with different potentials and different qualitative behavior of the Hubble parameter.

Given $U(\sigma)=\xi\sigma^2$, and taken $Y(\sigma)$ as a generic polynomial:

$$Y(\sigma)=\sum_{k=0}^{N}A_k\sigma^k, \tag{90}$$

where A_k are constants, from (25) one can obtain $F(\sigma)$ and show that it does not depend on A_1. For $N=2$, we obtain

$$F(\sigma)=\frac{4\xi\left((16\xi+1)A_0-(8\xi+1)A_2\sigma^2\right)\sigma}{(8\xi+1)(16\xi+1)}+\tilde{c}_0\sigma^{-1-1/(4\xi)}. \tag{91}$$

When $\tilde{c}_0=0$, $F(\sigma)$ is a cubic polynomial and Eq. (7) has the general solution which can be written in terms of elementary functions:

$$\sigma(t)=\pm\frac{\sqrt{(16\xi+1)^2A_0^2c_2e^{-\omega t}+(8\xi+1)(16\xi+1)A_2A_0}}{(16\xi+1)A_0c_2e^{-\omega t}+(8\xi+1)A_2}, \tag{92}$$

where $\omega=8\xi A_0/(8\xi+1)$, c_2 is an arbitrary integration constant and we assume that $\xi\neq-1/8$.

The corresponding potential, $V(\sigma)$, is the sixth degree polynomial which, for example, when $\xi=1$ has the following form:

$$V(\sigma)=\frac{910}{289}A_2^2\sigma^6+\frac{156}{17}A_1A_2\sigma^5+\left[6A_1^2+\frac{2236}{153}A_0A_2\right]\sigma^4+\frac{52}{3}A_0A_1\sigma^3+\frac{910}{81}A_0^2\sigma^2\,.$$

Let us analyze the cosmological consequences of the solution obtained. If $\omega>0$, then

$$\lim_{t\to\infty}\sigma(t)=\pm\frac{(16\xi+1)\sqrt{A_0}}{(8\xi+1)\sqrt{A_2}}\,. \tag{93}$$

In the case $\omega<0$, the function $\sigma(t)$ tends to zero at late times. Hence, the Hubble parameter tends to a constant at late times for any case[2]. On Fig. 2 we consider $H(t)$ for different values of A_1, for σ positive, $\xi=1$, $A_2=1$, and $A_0=2$. The solution $\sigma(t)$ in (92) is associated with different behaviors of the Hubble parameter. In particular, at $A_1=-4$ and $A_1=-2$ we get a non-monotonic behavior of $H(t)$. The behavior of the Hubble parameter, similar to the case $A_1=-4$, has been found in the quintom models with the sixth degree polynomial potential [21]. Such behavior can, in principle,

---

[2] We choose $Y(\sigma)$ to be a quadratic polynomial and therefore $A_2\neq0$.



Figure 2: The function $H(t)$, given by (90), with $A_1 = -6$, $A_1 = -4$, $A_1 = -2$, and $A_1 = 0$ (from left to right). At all pictures we use $A_2 = 1$, $A_0 = 2$, and $c_2 = 100000$.

describe both inflation and the late-times accelerating expansion of the Universe and is thus physically relevant.

In this section, we have obtained new induced gravity models, with a polynomial potential and physically interesting behavior of the Hubble parameter. Let us note that the explicit form of the Hubble parameter $H(t)$ is too complicated to be guessed and used in the reconstruction approach, proposed in [29]. We have obtained a three-parameter set of the induced gravity models with sixth degree polynomial potentials. The freedom of choice of parameters A_i allows one to set additional conditions on the model.

7 Construction of models with polynomial potentials

On considering cosmological models with scalar fields, which are inspired by some fundamental theory, for example, string field theory, it is difficult to get the exact form of the potential, but it is possible to get, at least, some properties of the potential. In [17, 21] string field theory inspired models with minimally coupled scalar fields and polynomial potentials have been considered. Let us address the question regarding what functions F and U are associated with a polynomial potential.

To get a polynomial potential from formula (11) we assume that $F(\sigma)$ is an arbitrary N degree polynomial:

$$F(\sigma) = \sum_{k=0}^{N} B_n \sigma^n. \tag{94}$$

If $U = \xi\sigma^2$, then from Eq. (8) we get

$$Y(\sigma) = \frac{B_0}{2}\left(1 + \frac{1}{4\xi}\right)\sigma^{-1} - \frac{B_2}{4\xi}(12\xi + 1)\sigma\ln(\frac{\sigma}{\sigma_0}) + Y_P(\sigma), \tag{95}$$

where $Y_P(\sigma)$ is a polynomial. The term proportional to σ^{-1} gives the constant term in the potential (11), thus only the term proportional to $\sigma\ln(\sigma)$ should be eliminated to obtain a polynomial potential. Hence, we come to the conclusion that for an arbitrary polynomial $F(\sigma)$ we get a polynomial potential if $\xi = -1/12$. This value of ξ corresponds to the case of conformal coupling [29]. For other values of ξ we get a polynomial potential only if $B_2 = 0$.

If U is an arbitrary quadratic polynomial, we get the following result: the coefficients of $Y(\sigma)$, proportional to B_0 and B_1 are always polynomial, those proportional to B_2 and B_3 are polynomial if $\xi = -1/12$. The coefficient proportional to B_4 includes logarithmic terms. The function $Y(\sigma)$ also contains the term, proportional to $\sqrt{U(\sigma)}$, multiplied by an integration constant, therefore, in order to get a polynomial $V(\sigma)$ this integration constant should be zero. Hence, to get a polynomial potential, we should either set $N = 1$ or $N = 3$ and $\xi = -1/12$. In the latter case, we obtain a sixth degree potential for nonzero B_3. For $B_3 = 0$ (or equivalently $N = 2$), the degree or the potential can not be more than four.

8 Conclusion

In this article we employed the reconstruction procedure for cosmological models with non-minimally coupled scalar fields evolving on a flat Friedmann-Robertson-Walker background. Besides their cosmological implications, models with non-minimally coupled scalar fields, including the Hilbert-Einstein term, are interesting because of their connection with particle physics. In particular we refer to models of inflation, where the role of the inflaton is played by the Higgs field non-minimally coupled to gravity. Such models have great relevance and were intensively discussed during the last years [49].

It is difficult to apply the reconstruction procedure proposed in [29] in the induced gravity context to more general modified gravity models. For these models, with the superpotential method, one can find many exact solutions for the homogeneous scalar field-gravity system and the corresponding potential of the scalar field. Such a method translates the problem of integrating a second order differential equation with time dependent coefficients into the integration of a first order differential equation once either the Hubble parameter is expressed as a function of the scalar field or the behavior of the scalar field itself is given. Let us note that both the variants of the superpotential technique we are referring to do not need the expression of the Hubble parameter in terms of the cosmic time or of the scale factor. This method has been actively employed in cosmological models with minimally coupled scalar fields [19, 21, 22, 23], but, to the best of our knowledge, it has not been applied to cosmological models with non-minimally scalar fields. Hence the superpotential method is a powerful tool to find exact analytical solutions of cosmological interest.

In this article we used this method to find exact analytical solutions leading to some physically interesting behaviors of the Hubble parameter. First we have found the potentials and the corresponding evolutions of the associated scalar field leading to de Sitter and power-law solutions. Power-law solutions reproduce the expansion of the Universe driven by a barotropic perfect fluid. For the former case we could formally reconstruct the dynamics of the scalar field-gravity system on assuming a generic coupling between the field and the Ricci scalar. In such a case both the superpotential technique and the reconstruction procedure proposed in [29] can be applied. We then specialized the results to the non-minimally coupled case with a general coupling constant ξ. The case of power-law expansion is more involved and we could only find and analyze some solutions with the parameters suitably tuned. Still, however, the reconstruction could only be performed easily by means of the superpotential technique.

We further investigated a few models having a de Sitter/anti de Sitter attractor far in the past or in the future. In the non-minimal coupling case the dynamics of a scalar field with a suitable quartic potential has been shown to have these attractors. In the induced gravity limit, we then also found a set of sixth degree polynomial potentials leading to a cosmic evolution with a constant $H(t)$ behavior both in the past and in the future.

Polynomial potential frequently appear as a result of the reconstruction procedure because polynomials are integrable and they can be systematically used for obtaining exact solutions with the superpotential technique. Hence we finally analyzed the sufficient conditions on the functions U and F to get a model with a polynomial potential.

Acknowledgements. We are grateful to A.A.Starobinsky for useful discussions. A.K. was partially supported by the RFBR grant 11-02-00643. Research of S.V. is supported in part by the RFBR grant

11-01-00894, by the Russian Ministry of Education and Science under grant NSh-3920.2012.2, and by contract CPAN10-PD12 (ICE, Barcelona, Spain).

References

[1] A. Riess et al, Astron. J. 116 (1998) 1009; S. J. Perlmutter et al, Astroph. J. 517 (199) 565.

[2] T. Padmanabhan, *Phys. Rept.* **380** (2003) 235–320, arXiv:hep-th/0212290;
P. Frampton, *Dark Energy – a Pedagogic Review*, arXiv:astro-ph/0409166;
E.J. Copeland, M. Sami, and Sh. Tsujikawa, *Int. J. Mod. Phys. D* **15** (2006) 1753–1936, arXiv:hep-th/0603057;
A.D. Dolgov, *Phys. Part. Nucl.* **43** (2012) 273–293;
K. Bamba, S. Capozziello, S. Nojiri, and S.D. Odintsov, Astrophys. Space Sci. 342 (2012) 155–228, arXiv:1205.3421

V. Sahni and A.A. Starobinsky, *Int. J. Mod. Phys.* D **9** (2000) 373–444, arXiv:astro-ph/9904398;
P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys.* **75** (2003) 559–606, arXiv:astro-ph/0207347;
V. Sahni, Class. Quantum Grav. **19** (2002) 3435–3448, arXiv:astro-ph/0202076 ;
V. Sahni and A.A. Starobinsky, *Int. J. Mod. Phys.* D **15** (2006) 2105, arXiv:astro-ph/0610026

[3] Y. Fujii and K. Maeda, *The Scalar–Tensor Theory of Gravitation*, Cambridge University Press, Cambridge, 2004

[4] S. Nojiri and S.D. Odintsov, *Int. J. Geom. Meth. Mod. Phys.* **4** (2007) 115–146, arXiv:hep-th/0601213;
S. Nojiri and S.D. Odintsov, *Phys. Rept.* **505** (2011) 59–144, arXiv:1011.0544

[5] S. Capozziello and V. Faraoni, *Beyond Einstein Gravity: A Survey of Gravitational Theories for Cosmology and Astrophysics*, Fund. Theor. Phys. **170**, Springer, New York, 2011

[6] A. de Felice and Sh. Tsujikawa, *Living Rev. Rel.* **13** (2010) 3, arXiv:1002.4928

[7] A. A. Starobinsky, Lect. Notes in Phys. **246** (1986) 107;
A. D. Linde, *Particle Physics and Inflationary Cosmology*, Chur, Switzerland: Harwood, 1990.

[8] J.E. Lidsey, A.R. Liddle, E.W. Kolb, E.J. Copeland, T. Barreiro, M. Abney, *Rev. Mod. Phys.* **69** (1997) 373–410, arXiv:astro-ph/9508078;
C.M. Peterson, M. Tegmark, *Phys. Rev.* D **83** (2011) 023522 , arXiv:1005.4056;
Shi Pi, M. Sasaki, *J. Cosmol. Astropart. Phys.* **1210** (2012) 051, arXiv:1205.0161

[9] E. Elizalde, Sh. Nojiri, S.D. Odintsov, D. Saéz-Gómez, V. Faraoni, *Phys. Rev.* D **77** (2008) 106005, arXiv:0803.1311

[10] A.A. Starobinsky, *JETP Lett.* **68** (1998) 757–763, arXiv:astro-ph/9810431

[11] A.B. Burd and J.D. Barrow, *Nucl. Phys.* B **308** (1988) 929,

[12] J.D. Barrow, *Phys. Lett.* B **235** (1990) 40.

[13] A. Yurov, *Phantom scalar fields result in inflation rather than Big Rip*, arXiv:astro-ph/0305019

[14] Z.K. Guo, N. Ohta and Y.Z. Zhang, *Mod. Phys. Lett.* A **22** (2007) 883, arXiv:astro-ph/0603109

[15] Z.K. Guo, N. Ohta and Y.Z. Zhang, *Phys. Rev.* D **72** (2005) 023504, arXiv:astro-ph/0505253

[16] T.D. Saini, S. Raychaudhury, V. Saini, and A.A. Starobinsky, *Phys. Rev. Lett.* **85** (2000) 1162–1165, arXiv:astro-ph/9910231

[17] I.Ya. Aref'eva, A.S. Koshelev, and S.Yu. Vernov, *Theor. Math. Phys.* **148** (2006) 895–909, arXiv:astro-ph/0412619

[18] S. Capozziello, S. Nojiri, and S.D. Odintsov, *Phys. Lett.* B **634** (2006) 93–100, arXiv:hep-th/0512118

[19] A.G. Muslimov, *Class. Quant. Grav.* **7** (1990) 231–237,
D.S. Salopek and J.R. Bond, *Phys. Rev.* D **42** (1990) 3936–3962,
V.M. Zhuravlev, S.V. Chervon and V.K. Shchigolev, J. Exp. Theor. Phys. **87** (1998) 223;
D. Bazeia, C.B. Gomes, L. Losano, and R. Menezes, *Phys. Lett.* B **633** (2006) 415–419, arXiv:astro-ph/0512197;
P.K. Townsend, *Class. Quant. Grav.* **25** (2008) 045017, arXiv:0710.5178;
A.V. Yurov, V.A. Yurov, S.V. Chervon, and M. Sami, *Theor. Math. Phys.* **166** (2011) 259–269;
A.Yu. Kamenshchik and S. Manti, *Gen. Rel. Grav.* **44** (2012) 2205–2214, arXiv:1111.5183;
H.-Ch. Kim, *Exact solutions in Einstein cosmology with a scalar field*, arXiv:1211.0604

[20] S. Nojiri and S.D. Odintsov, *J. Phys. Conf. Ser.* **66** (2007) 012005, arXiv:hep-th/0611071

[21] I.Ya. Aref'eva, A.S. Koshelev, and S.Yu. Vernov, *Phys. Rev.* D **72** (2005) 064017, arXiv:astro-ph/0507067;
S.Yu. Vernov, *Theor. Math. Phys.* **155** (2008) 544–556, arXiv:astro-ph/0612487

[22] A.A. Andrianov, F. Cannata, A.Yu. Kamenshchik, and D. Regoli, *J. Cosmol. Astropart. Phys.* **0802** (2008) 015, arXiv:0711.4300

[23] I.Ya. Aref'eva, N.V. Bulatov, and S.Yu. Vernov, *Theor. Math. Phys.* **163** (2010) 788–803, arXiv:0911.5105

[24] T. Padmanabhan, *Phys. Rev.* D **66** (2002) 021301

[25] A. Feinstein, *Phys. Rev.* D **66** (2002) 063511

[26] V. Gorini, A.Y. Kamenshchik, U. Moschella, and V. Pasquier, *Phys. Rev.* D **69** (2004) 123512, arXiv:hep-th/0311111

[27] V.K. Shchigolev and M.P. Rotova, *Mod. Phys. Lett.* A **27** (2012) 1250086, arXiv:1203.5030

[28] B. Boisseau, G. Esposito-Farese, D. Polarski, and A.A. Starobinsky, *Phys. Rev. Lett.* **85** (2000) 2236, arXiv:gr-qc/0001066

[29] A.Yu. Kamenshchik, A. Tronconi, and G. Venturi, *Phys. Lett.* B **702** (2011) 191–196, arXiv:1104.2125

[30] Taotao Qiu, *J. Cosmol. Astropart. Phys.* **1206** (2012) 041, arXiv:1204.0189

[31] K. Bamba, S. Nojiri, and S.D. Odintsov, *Phys. Rev.* D **77** (2008) 123532, arXiv:0803.3384;
E. Elizalde and A.J. López-Revelles, *Phys. Rev.* D **82** (2010) 063504, arXiv:1004.5021;
E. Elizalde, A.J. López-Revelles, S.D. Odintsov, and S.Yu. Vernov, Cosmological models with Yang–Mills fields, arXiv:1201.4302

[32] P.K.S. Dunsby, E. Elizalde, R. Goswami, S. Odintsov, and D. Saez-Gomez, *Phys. Rev.* D **82** (2010) 023519, arXiv:1005.2205;
R. Myrzakulov, D. Saez-Gomez, and A. Tureanu, *Gen. Rel. Grav.* **43** (2011) 1671–1684, arXiv:1009.0902

[33] M. Jamil, D. Momeni, M. Raza, and R. Myrzakulov, *Eur. Phys. J.* C **72** (2012) 1999, arXiv:1107.5807;
K. Bamba, R. Myrzakulov, S. Nojiri, and S.D. Odintsov, *Phys. Rev.* D **85** (2012) 104036, arXiv:1202.4057

[34] C. Deffayet and R.P. Woodard, *J. Cosmol. Astropart. Phys.* **0908** (2009) 023, arXiv:0904.0961

[35] T.S. Koivisto, *Phys. Rev.* D **77** (2008) 123513, arXiv:0803.3399;
E. Elizalde, E.O. Pozdeeva, and S.Yu. Vernov, *Phys. Rev.* D **85** (2012) 044002, arXiv:1110.5806;
S.Yu. Vernov, *Phys. Part. Nucl.* **43** (2012) 694–696, arXiv:1202.1172;
E. Elizalde, E.O. Pozdeeva, and S.Yu. Vernov, *Reconstruction Procedure in Nonlocal Models*, arXiv:1209.5957

[36] F. Cooper and G. Venturi, *Phys. Rev.* D **24** (1981) 3338.

[37] F. Finelli, A. Tronconi and G. Venturi, *Phys. Lett.* B **659** (2008) 466.

[38] A. Cerioni, F. Finelli, A. Tronconi and G. Venturi, *Phys. Lett.* B **681** (2009) 383.

[39] A. Cerioni, F. Finelli, A. Tronconi and G. Venturi, *Phys. Rev.* D **81** (2010) 123505.

[40] A. Tronconi and G. Venturi, Phys. Rev. D **84** (2011) 063517 [arXiv:1011.3958 [gr-qc]].

[41] A. Y. Kamenshchik, A. Tronconi and G. Venturi, Phys. Lett. B **713** (2012) 358 [arXiv:1204.2625 [gr-qc]].

[42] M. Sami, M. Shahalam, M. Skugoreva, A. Toporensky, arXiv:1207.6691 [hep-th] [arXiv:1207.6691 [hep-th]].

[43] J.L. Cervantes-Cota, R. de Putter and E.V. Linder, *J. Cosmol. Astropart. Phys.* **1012** (2010) 019

[44] B. L. Spokoiny, Phys. Lett. B **147**, 39 (1984);
T. Futamase and K. -i. Maeda, Phys. Rev. D **39**, 399 (1989);
D. S. Salopek, J. R. Bond and J. M. Bardeen, Phys. Rev. D **40**, 1753 (1989);
R. Fakir and W. G. Unruh, Phys. Rev. D **41**, 1783 (1990).

[45] A. O. Barvinsky and A. Y. Kamenshchik, Phys. Lett. B **332**, 270 (1994);
A. O. Barvinsky and A. Y. Kamenshchik, Nucl. Phys. B **532**, 339 (1998);
A. O. Barvinsky, A. Y. Kamenshchik, C. Kiefer and C. F. Steinwachs, Phys. Rev. D **81**, 043530 (2010).

[46] A. Brandhuber and K. Sfetsos, *J. High Energy Phys.* **9910** (1999) 013, hep-th/9908116;
O. DeWolfe, D.Z. Freedman, S.S. Gubser, A. Karch, *Phys. Rev.* D **62** (2000) 046008; hep-th/9909134;
A.S. Mikhailov, Yu.S. Mikhailov, M.N. Smolyakov, I.P. Volobuev, *Class. Quantum Grav.* **24** (2007) 231–242, hep-th/0602143;
K. Skenderis and P.K. Townsend, *Phys. Rev.* D **74** (2006) 125008, arXiv:hep-th/0609056

[47] T. Biswas, A. Mazumdar, and W. Siegel, *J. Cosmol. Astropart. Phys.* **0603** (2006) 009, arXiv:hep-th/0508194;
I.Ya. Aref'eva, L.V. Joukovskaya, and S.Yu. Vernov, *J. Phys. A: Math. Theor.* **41** (2008) 304003, arXiv:0711.1364;
T. Biswas, T. Koivisto, and A. Mazumdar, *J. Cosmol. Astropart. Phys.* **1011** (2010) 008, arXiv:1005.0590;
T. Biswas, A.S. Koshelev, A. Mazumdar, and S.Yu. Vernov, *J. Cosmol. Astropart. Phys.* **1208** (2012) 024, arXiv:1206.6374

[48] R. Gannouji, D. Polarski, A. Ranquet and A. A. Starobinsky, *J. Cosmol. Astropart. Phys.* **0609** (2006) 016, astro-ph/0606287.

[49] F. L. Bezrukov and M. Shaposhnikov, Phys. Lett. B **659**, 703 (2008);
A. O. Barvinsky, A. Y. Kamenshchik and A. A. Starobinsky, JCAP **0811**, 021 (2008);
F. L. Bezrukov, A. Magnin and M. Shaposhnikov, Phys. Lett. B **675**, 88 (2009);
A. De Simone, M. P. Hertzberg and F. Wilczek, Phys. Lett. B **678**, 1 (2009);
F. Bezrukov and M. Shaposhnikov, JHEP **0907**, 089 (2009);
A. O. Barvinsky, A. Y. Kamenshchik, C. Kiefer, A. A. Starobinsky and C. Steinwachs, JCAP **0912**, 003 (2009);
F. Bezrukov, A. Magnin, M. Shaposhnikov and S. Sibiryakov, JHEP **1101**, 016 (2011);
A. O. Barvinsky, A. Y. Kamenshchik, C. Kiefer, A. A. Starobinsky and C.F. Steinwachs, Eur.Phys. J. C **72**, 2219 (2012).